



## **DRAFT REPORT**

# **MUTAGENICITY STUDY OF PROSOLUTION TABLETS AS PER OECD GUIDELINE NO. 471 - BY SALMONELLA TYPHIMURIUM REVERSE MUTATION ASSAY**

**STUDY NO: VLTO-100219**

**Study Completion Date: xx.xx. 2010**

### **SPONSOR**

#### **DM CONTACT MANAGEMENT.**

100-645 TYEE ROAD VICTORIA  
BC V9A6X5, CANADA.

### **TEST FACILITY**

#### **VEDIC LIFESCIENCES PVT. LTD.**

203, MORYA LANDMARK-I,  
OFF LINK ROAD, ANDHERI (W),  
MUMBAI – 400 053  
INDIA



## **STATEMENT OF COMPLIANCE**

**To the best of our knowledge and belief, this Study entitled “Mutagenicity study of Prosolution tablets by Salmonella typhimurium, Reverse Mutation Assay” was performed under my supervision in compliance with the test guidelines laid down in OECD – 471”. The objectives laid down in the study protocol were achieved.**

**No unforeseen circumstances were observed which might have affected the quality or integrity of the study.**

**Jayesh Chaudhary  
CEO, Vedic Lifesciences Pvt. Ltd.**

**Deepali Jadhav  
Executive**



## **CERTIFICATE**

We certify that the work reported here is a true and authentic report of the study entitled, “Mutagenicity study of Prosolution tablets by Salmonella typhimurium, Reverse Mutation Assay as per the OECD guidelines – 471”, based on the experiment conducted in one of the partnered Toxicology Laboratory Services of VEDIC LIFESCIENCES PVT LTD (B-203 Morya Landmark I, Off New Link Road, Andheri (W), Mumbai - 400 053,) India. The results presented here are faithful reflection of data collected during the study.



## **TABLE OF CONTENTS**

<b>SL. NO.</b>	<b>TITLE</b>	<b>PAGE NO.</b>
	QUALITY ASSURANCE STATEMENT -----	6
	STATEMENT OF CONFIDENTIALITY -----	6
	DECLARATION -----	7
1.	STUDY DETAILS -----	8
2.	MONITORING PERSONNEL -----	9
3.	SUMMARY -----	10
4.	OBJECTIVE -----	11
5.	MATERIALS AND METHODS -----	11
	5.1 TEST ARTICLES -----	11
	5.2 TEST SYSTEM -----	11
	5.3 GENETIC CHARACTERIZATION OF TESTER STRAINS -----	12
	5.4 METABOLIC ACTIVATION (S9) DETAILS -----	12
	5.5 STERILITY TEST OF THE TEST ITEM -----	13
	5.6 SOLUBILITY TEST OF THE TEST ITEM -----	13
	5.7 TEST ITEM PREPARATION -----	13
	5.8 PRECIPITATION TEST -----	13
	5.9 DURATION OF CULTURE -----	13
	5.10 INITIAL CYTOTOXICITY TEST -----	13
6.	PLATE INCORPORATION METHOD FOR THE MUTAGENICITY ASSAY -----	14
7.	OBSERVATIONS -----	15
8.	DATA ANALYSIS -----	15
9.	ARCHIVES -----	15
10.	RESULTS AND DISCUSSION -----	15
	10.1 GENETIC CHARACTERIZATION OF TESTER STRAINS -----	15
	10.2 METABOLIC ACTIVATION -----	15
	10.3 STERILITY TEST -----	16
	10.4 SOLUBILITY TEST -----	16
	10.5 PRECIPITATION TEST -----	16
	10.6 INITIAL CYTOTOXICITY TEST -----	16
	10.7 PLATE INCORPORATION METHOD FOR THE MUTAGENICITY ASSAY -----	17
11.	INTERPRETATION -----	18
12.	CONCLUSION -----	18



## **Summary Tables**

Table 1: Summary of Initial Cytotoxicity Test .....	19
Table 2: Summary of Colony Counts of Revertants of Trial - I .....	20
Table 3: Summary of Colony Counts of Revertants of Trial - II .....	21

## **Appendices**

Appendix 1: Individual Plate Colony Counts of Initial Cytotoxicity Test .....	22
Appendix 2: Individual Plate Colony Counts of Revertants of Trial - I .....	23
Appendix 3: Individual Plate Colony Counts of Revertants of Trial - II .....	24
Appendix 4: Sterility Status of the Study .....	25



## **QUALITY ASSURANCE STATEMENT**

The study entitled “Mutagenicity study of Prosolution tablets by Salmonella typhimurium, Reverse Mutation Assay” has been inspected in the spirit of OECD Guidelines 471

This study was inspected and findings reported to Management and to the Study Director.

Inspections were performed according to the Standard Operating Procedures of the Quality Assurance Unit. The report was audited against the approved study plan and pertinent raw data and accurately reflects the raw data.

## **STATEMENT OF CONFIDENTIALITY**

This report which contains **CONFIDENTIAL** and **PROPRIETARY** information of **DM Contact Management**. will not be disclosed to anyone except the employees of this company wherever necessary or to persons authorized by law or judicial judgment without the expressed or written approval of Sponsor.



## **DECLARATION**

The Study Director hereby declares that the work was performed under his supervision and in accordance with the described procedures. It is assured that the reported results faithfully represent the raw data obtained during the experimental work. No circumstances have been left unreported which may have affected the quality or integrity of the data or which might have a potential bearing on the validity and reproducibility of this study.

The Study Director accepts overall responsibility for the technical conduct of the study as well as the interpretation, analysis, documentation and reporting of the results.



## 1. STUDY DETAILS

**1.1 TITLE** : Mutagenicity study of Prosolution tablets by Salmonella Typhimurium Reverse Mutation Assay (Ames test)

**1.2 STUDY NUMBER** : VLTO – 100219

**1.3 TESTING FACILITY** : VEDIC LIFESCIENCES PVT. LTD,  
203, MORYA LANDMARK-I,  
OFF LINK ROAD, ANDHERI (W),  
MUMBAI – 400 053, INDIA

**1.4 SPONSOR** : DM CONTACT MANAGEMENT  
100-645 TYEE ROAD  
VICTORIA, BC V9A6X5  
CANADA

**1.5 STUDY SCHEDULE** :

Study Initiation date : 26.04.2010

Metabolic activation –S9 preparation : 27.04.2010 to 05.05.2010

Solubility and precipitation test : 06.05.2010

Initial cytotoxicity test and colony counting : 08.05.2010 to 10.05.2010

Trial 1: test item exposure : 24.05.2010  
(With or without S9 activation)

Trial 1 : Colony counting : 26.05.2010  
(With or without S9 activation)

Trial 2 : Test item exposure : 28.05.2010  
(With or without S9 activation)

Trial 2 : Colony counting : 30.05.2010  
(with and without S9 activation)





## 2. MONITORING PERSONNEL

<b>Sr. No.</b>	<b>Responsibility</b>	<b>Personnel</b>	<b>Signature with date</b>
1.	MONITORING SCIENTIST	DEEPALI BHOSALE VEDIC LIFESCIENCES PVT.LTD MUMBAI	
2.	MANAGEMENT NOMINEE	JAYESH CHAUDHARY VEDIC LIFESCIENCES PVT.LTD MUMBAI	



### 3. SUMMARY

The test item Prosolution tablets manufactured and supplied by DM contact management, Canada was assessed for its mutagenic effects using Salmonella typhimurium. The study was conducted using Salmonella typhimurium strains: TA98, TA100, TA102, TA1535, and TA1537. The test item was tested at the concentration of 10, 50, 100, 500 and 1000 µg/plate using dimethyl sulphoxide (DMSO) as solvent based on the initial cytotoxicity test. The study was conducted, with and without the metabolic activation (S9) prepared from Aroclor 1254 induced in rat liver. The control, solvent control and appropriate positive controls (2-nitrofluorene, sodium azide and 9-aminoacridine, mitomycin C for trials “without metabolic activator” and 2-aminoanthracene for trials “with metabolic activator”) were tested simultaneously. Two trials were carried out for this study in triplicates. Data were statistically analyzed and expressed as mean ± SD.

From the experimental results obtained, the mean numbers of revertant colonies in the above mentioned concentrations were comparable to those of the control and solvent control, in both the trials, in the presence and absence of metabolic activation. There was no significant increase in number of revertant colonies at all concentrations tested. The number of revertant colonies in the positive controls has shown 3.0 to 23.5 fold increase under identical conditions.

#### Conclusion

From the results obtained, the test item Prosolution tablets is found to be non-mutagenic at the highest concentration 1000 µg/plate in Bacterial Reverse Mutation Test in the tester strain Salmonella typhimurium strains: TA98, TA100, TA102, TA1535, and TA1537.



#### 4. OBJECTIVE

To assess the potential of test item that can induce point gene mutations, viz., substitution, addition or deletion of one or a few DNA base pairs in the Salmonella typhimurium mutation. The experiment was performed by plate incorporation method.

#### 5. MATERIALS AND METHODS

##### 5.1 TEST ARTICLES

The following information was provided about the test article.

Test article	: Prosolution tablets
Characteristics	: White coloured tablets.
Batch No.	: NA
Date of Manufacture	: NA
Date of Expiry	: NA
Purity	: NA
Sponsor	: DM CONTACT MANAGEMENT 100-645 TYEE ROAD VICTORIA, BC V9A6X5 CANADA

##### 5.2 TEST SYSTEM

Tester Bacteria	: Salmonella typhimurium
Tester Strain Used	: TA 1537, TA 98.
Base substitution	: TA 100, TA 102 and TA 1535.
Source	: Moleular Toxicology, Inc.,157 ,Industrial park, Dr. Biine, NC 28607,USA



### 5.3 Genetic Characterization of Tester Strains

Strains were maintained as master plates and periodically checked for viable counts and genetic characteristics as mentioned below:

- a. Histidine requirement
- b. rfa mutation
- c. uvr B mutation
- d. R-factor (pKM 101 plasmid)
- e. Spontaneous reversion

### 5.4 Metabolic Activation (S9) Details

#### Preparation of S9 homogenate (Metabolic activator)

Liver microsomal enzymes were prepared from male Wistar rats induced with single exposure of Aroclor 1254 at a concentration of 500mg/kg body weight. After 5 days of Aroclor dosing, animals were fasted over night and sacrificed on 6<sup>th</sup> day. Livers were excised under aseptic condition. The preparation of the microsomal enzyme fraction was carried out with sterile glassware and solutions under ice cold condition. The liver was suspended in 3 volumes of 0.15 M KCl (3 ml/g of liver) and minced with sterile scissors. The minced liver was homogenized in a tissue homogenizer and was centrifuged at 9000 x g for 10 minutes in a refrigerated centrifuge. The supernatant was decanted and aliquots transferred into cryovials which were frozen and stored at -70°C.

#### Details of S9 homogenate

Date of preparation: 02.05.2010

Date of characterization: Sterility check: 02.05.2010; Activity check: 03.05.2010

#### Sterility check for S-9 fraction

Sterility was checked by streaking the supernatant fluid on nutrient agar plates and incubated at 37°C for 24 hours.



### **Activity check for S-9 homogenate**

Activity of the S-9 homogenate was determined using a pro-mutagen 2-aminoanthracene and Benzo(a)pyrene with the tester strain TA100.

### **5.5 Sterility Test of the Test item**

Test item was checked for its microbial load by pour plate technique

### **5.6 Solubility Test of the Test item**

The solubility test of test item was carried out with water and DMSO.

### **5.7 Test Item Preparation**

Test item stock solutions preparation: 1.25 g of the test substance was suspended in DMSO and volume was made up to 25 ml corresponding to 50 mg/ml stock.

### **5.8 Precipitation Test**

The test item was dissolved in DMSO and serially diluted to get concentrations of 1000, 2000, 3000, 4000 and 5000 µg/ml and precipitation test was carried out for the same.

### **5.9 Duration of Culture**

Cultures from genotyped master plates of each strain were grown in oxid nutrient broth No. 2 for 16-18 hours at 37°C ± 1°C in the shaking water bath set at approximately 68 shakes / minute and were assessed using viable count analysis.

### **5.10 Initial Cytotoxicity Test**

Based on the results of the precipitation test, the following concentrations were selected for the initial cytotoxicity test (µg/plate):

- a. 1000    b. 2000    c. 3000    d. 4000    e. 5000

Initial toxicity test was conducted using overnight TA100 tester strain, both in the presence and the absence of metabolic activator, in triplicate, along with concurrent solvent control (DMSO). The test item was judged as toxic based on decrease in the number of revertants/plate and/or the bacterial background lawn intensity.



## 6.0 PLATE INCORPORATION METHOD FOR THE MUTAGENICITY ASSAY

### Test System Identification:

Based on the initial cytotoxicity test results, the following concentrations were selected for the main study ( $\mu\text{g}/\text{plate}$ ):

a. 10    b. 50        c. 100    d. 500    e. 1000

The plates were labeled with the type of study number, strain, metabolic activation (with or without), trial number and concentration of test item.

### Plating:

Five concentrations of the test item were plated, with each of the following tester strains: TA98, TA100, TA102, TA1535 and TA1537 with and without metabolic activator. Two trials were conducted with three replicates per strain for each concentration of test item, both in the presence and absence of the metabolic activation.



## **7.0 OBSERVATIONS**

### **Effect on bacterial background lawn:**

The condition of the bacterial background lawn was evaluated for evidence of the test item toxicity using the code system.

### **Number of revertants:**

Revertant colonies for a given strain within the test item dilution series were counted manually.

## **8.0 DATA ANALYSIS**

Data was analyzed for differences among control, solvent control and positive control groups using ANOVA. Differences between the control, solvent control and treatment groups were tested by Dunnett's test at a 5% level ( $p \leq 0.05$ ) of significance.

## **9.0 ARCHIVING**

All test article, reference sample, study protocol, raw data and other documents generated during the course of this study together with a copy of final report will be stored in the archives of Vedic Lifesciences Pvt. Ltd., India for a period of one year from the date of submission of final report.

## **10.0 . RESULTS AND DISCUSSION**

### **10.1 Genetic Characterization of Tester Strains**

All the tester strains fulfilled the quality check criteria.

### **10.2 Metabolic activation**

#### **Sterility check for S-9 fraction:**

Sterility was checked by streaking the supernatant fluid on nutrient agar plates and incubated at 37°C for 24 hours. It was found sterile.



### **Activity check for S-9 homogenate:**

Activity of the S-9 homogenate was determined using a pro-mutagen 2-aminoanthracene and Benzo(a)pyrene with the tester strain TA100 and it was found to be active.

### **10.3 Sterility Test**

The test item was found to be sterile at tested conditions.

### **10.4 Solubility Test**

The test item was found to be suspended in DMSO at 50 mg/ml concentration.

Solvent : DMSO  
Batch No : R237F09  
Make : Rankem  
Storage : Room temperature

### **10.5 Precipitation Test**

It was found that the test item did not cause any precipitation at the concentration 5000 µg/plate in Minimal Glucose agar.

### **10.6 Initial Cytotoxicity Test**

In the test, with and without S9 activation, at the tested concentrations of 1000, 2000, 3000, 4000 and 5000 µg/plate the mean number of revertant colonies and the bacterial background lawn were compared to that of the solvent control. The test item was found to be cytotoxic at and above 2000 µg/plate with decrease in number of revertant colonies and depletion in bacterial background lawn as compared to that of the solvent control. Refer Table 1, Appendix 1





## 10.7 Plate Incorporation Method for the Mutagenicity Assay

### **Trial I:**

The concentrations tested at 10, 50, 100, 500 and 1000 µg/plate showed very close resemblance to the solvent control DMSO in both with and without metabolic activation. There was no significant increase in number of revertant colonies and no change in bacterial background lawn as compared to that of the solvent control, among the tester strains. The specific positive controls tested simultaneously produced approximately 3.0 to 23.5 fold increase in mean number of revertants as compared to the solvent control. Refer Table 2, Appendix 2.

### **Trial II:**

Similar results were observed in the second trial, the concentrations tested at 10, 50, 100, 500 and 1000 µg/plate showed very close resemblance to the solvent control DMSO in both with and without metabolic activation. There was no significant increase in number of revertant colonies and no change in bacterial background lawn as compared to that of the solvent control, among the tester strains. The mean number of revertant colonies/plate and bacterial background lawn in the solvent control for the tested strains were comparable to that of solvent control. The specific positive controls tested simultaneously produced approximately 3.0 to 22.5 fold increase in the mean number of revertants as compared to the solvent control. Refer Table 3, Appendix 3.

Statistical analysis of the combined results from both trials indicated no significance in all the five concentrations tested which was compared to the respective solvent control in any of the five *Salmonella* strains.



## 11.0 INTERPRETATION

The test item Prosolution tablets was assayed for Bacterial Reverse Mutation Test at the concentration of 10, 50, 100, 500, 1000 µg/plate using *Salmonella typhimurium* strains; TA98, TA100, TA102, TA1535 and TA1537. In the two trials conducted, with metabolic activation (S9) and without metabolic activation, the number of revertant colonies in the five different test item concentrations did not increase significantly over the solvent control, while, the positive controls tested simultaneously showed a 3.0 to 23.5 fold increase in the number of revertant colonies/plate. This was observed for all the five tester strains.

## 12.0 CONCLUSION

From the results obtained, the test item Prosolution tablets is found to be non-mutagenic at the highest concentration 1000µg/plate in Bacterial Reverse Mutation Test in the tester strain *Salmonella typhimurium* strains: TA98, TA100, TA102, TA1535, and TA1537.



**TABLE – 1: SUMMARY OF INITIAL CYTOTOXICITY TEST**

Treatment		Control		Vehicle Control		Test item (µg/plate)									
Test concentration (µg/plate)		100µl of autoclaved distilled water		100µl of DMSO		1000		2000		3000		4000		5000	
--		Mean ± SD	Lawn Intensity	Mean ± SD	Lawn Intensity	Mean ± SD	Lawn Intensity	Mean ± SD	Lawn Intensity	Mean ± SD	Lawn Intensity	Mean ± SD	Lawn Intensity	Mean ± SD	Lawn Intensity
No. of Revertants / Plate Salmonella typhimurium TA 100	With S-9	99.7 ±0.6	4+	99.7 ±1.5	4+	99.7 ±1.2	4+	94.7 ±1.5	4+	85.7 ±3.1	4+	85.0 ±3.6	3+	82.3 ±1.5	3+
	Without S-9	99.7 ±1.5	4+	100.7 ±1.2	4+	100.3 ±1.2	4+	96.3 ±2.1	4+	86.7 ±2.5	4+	82.3 ±2.5	3+	84.3 ±3.1	3+

Lawn intensity:

4+= Thick lawn: Distinguished by a healthy (Normal) background lawn comparable to solvent control plates.

3+= Slightly thin lawn: Distinguished by a noticeable lawn (Slightly thinning of the background lawn reduced) compared to solvent control plates.



**TABLE – 2: SUMMARY OF COLONY COUNTS OF REVERTANTS OF TRIAL – I**

Treatment	Test concentration (µg/plate)	No. of Revertants (Mean of 3 plates)									
		With S-9					Without S-9				
		TA 98	TA 100	TA 102	TA 1535	TA 1537	TA 98	TA 100	TA 102	TA 1535	TA 1537
Control	100µl of autoclaved distilled water	17.7 ±1.5	101.0 ±1.0	248.3 ±10.8	20.0 ±1.0	8.3 ±0.6	17.0 ±1.0	100.0 ±1.0	253.3 ±3.1	18.0 ±2.0	6.0 ±1.0
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Vehicle Control	100µl of DMSO	17.7 ±0.6	100.7 ±1.5	247.3 ±10.3	20.3 ±1.5	7.0 ±2.0	15.7 ±1.5	100.7 ±1.2	256.0 ±3.6	18.0 ±1.0	7.7 ±1.5
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Test Item	50	17.3 ±1.5	101.3 ±1.5	251.7 ±1.5	20.0 ±1.0	7.0 ±1.0	15.0 ±1.0	100.7 ±0.6	256.7 ±2.1	18.7 ±1.2	7.0 ±1.7
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	0.9
	100	16.3 ±0.6	101.0 ±1.0	251.0 ±1.0	19.7 ±1.5	6.7 ±1.2	15.0 ±1.0	101.0 ±1.7	257.7 ±2.5	18.7 ±2.5	8.0 ±1.0
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	0.9	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
	500	18.0 ±1.0	99.3 ±1.2	252.0 ±3.6	18.3 ±2.5	7.3 ±2.1	14.7 ±0.6	101.0 ±1.0	257.0 ±2.6	17.7 ±1.5	7.3 ±1.2
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	1.0	1.0	1.0	0.9	1.0	0.9	1.0	1.0	1.0	1.0
	1000	16.7 ±0.6	100.0 ±1.0	249.3 ±8.1	17.3 ±1.5	7.0 ±1.0	15.0 ±0.0	101.3 ±0.6	257.3 ±2.5	17.7 ±2.1	6.7 ±2.1
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	0.9	1.0	1.0	0.9	1.0	1.0	1.0	1.0	1.0	0.9
	2000	18.3 ±0.6	100.0 ±1.0	247.0 ±6.1	17.3 ±0.6	6.0 ±1.4	15.0 ±1.0	100.7 ±0.6	256.0 ±2.6	18.7 ±0.6	6.7 ±0.6
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	1.0	1.0	1.0	0.9	0.9	1.0	1.0	1.0	1.0	0.9
	Positive Control	100µl	195.7 ±0.6	321.0 ±1.0	754.7 ±16.2	190.7 ±6.4	120.3 ±0.6	213.7 ±1.5	313.0 ±2.0	770.3 ±13.4	195.3 ± 3.2
Lawn Intensity		4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Fold increase		11.1	3.2	3.1	9.4	17.2	13.6	3.1	3.0	10.9	22.6

Values of Revertants are in Mean±SD



**TABLE – 3: SUMMARY OF COLONY COUNTS OF REVERTANTS OF TRIAL - II**

Treatment	Test concentration (µg/plate)	No. of Revertants (Mean of 3 plates)									
		With S-9					Without S-9				
		TA 98	TA 100	TA 102	TA 1535	TA 1537	TA 98	TA 100	TA 102	TA 1535	TA 1537
Control	100µl of autoclaved distilled water	17.3 ±1.5	98.7 ±0.6	250.7 ±3.1	17.0 ±1.0	7.0 ±1.0	17.7 ±1.5	99.7 ±0.6	253.3 ±3.1	17.0 ±1.0	8.7 ±1.2
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Vehicle Control	100µl of DMSO	18.7 ±0.6	98.7 ±1.2	257.3 ±2.5	17.3 ±2.1	6.0 ±1.0	19.3 ±0.6	99.3 ±1.5	256.0 ±2.6	17.3 ±1.2	9.0 ±1.0
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Test Item	50	18.3 ±2.1	100.0 ±1.0	257.7 ±1.5	18.0 ±1.7	6.7 ±2.1	19.3 ±1.5	98.7 ±1.2	255.7 ±1.5	18.0 ±1.7	9.3 ±1.2
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	1.0	1.0	1.0	1.0	1.1	1.0	1.0	1.0	1.0	1.0
	100	17.0 ±1.0	101.0 ±1.0	255.7 ±1.5	17.7 ±2.9	6.3 ±1.5	17.0 ±1.0	99.7 ±1.5	257.3 ±2.5	18.3 ±1.5	9.0 ±1.0
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	0.9	1.0	1.0	1.0	1.1	0.9	1.0	1.0	1.1	1.0
	500	17.3 ±1.2	99.3 ±0.6	256.7 ±4.2	16.7 ±1.5	6.3 ±0.6	17.3 ±1.5	100.3 ±1.5	255.7 ±3.2	18.7 ±2.1	8.0 ±1.0
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	0.9	1.0	1.0	1.0	1.1	0.9	1.0	1.0	1.1	0.9
	1000	17.0 ±1.0	99.7 ±1.5	249.7 ±1.5	17.7 ±1.5	6.0 ±1.0	16.7 ±0.6	98.7 ±0.6	257.7 ±2.5	17.0 ±1.0	7.3 ±1.2
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	0.9	1.0	1.0	1.0	1.0	0.9	1.0	1.0	1.0	0.8
	2000	16.7 ±0.6	97.7 ±1.5	252.3 ±3.1	16.7 ±1.5	6.3 ±1.5	16.7 ±0.6	99.7 ±1.5	255.0 ±2.6	16.7 ±0.6	8.0 ±1.0
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	Fold increase	0.9	1.0	1.0	1.0	1.1	0.9	1.0	1.0	1.0	0.9
	Positive Control	100µl	195.7 ±3.5	321.7 ±3.8	764.7 ±4.7	198.3 ±6.5	118.0 ±2.6	217.7 ±2.5	324.0 ±2.6	770.0 ±2.6	200.0 ±5.0
Lawn Intensity		4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Fold increase		10.5	3.3	3.0	11.4	19.7	11.3	3.3	3.0	11.5	21.8

Values of Revertanats are in Mean±SD



**APPENDIX – 1: INDIVIDUAL PLATE COLONY COUNTS OF INITIAL CYTOTOXICITY TEST**

Treatment		Control			Vehicle Control			Test item (µg/plate)														
Test concentration (µg/plate)		100µl of autoclaved distilled water			100µl of DMSO			1000			2000			3000			4000			5000		
Plate No.		1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3	1	2	3
No. of Revertants/Plate <i>Salmonella Typhimurium</i> TA 100	With S-9	99	100	100	101	98	100	99	101	99	95	96	93	83	85	89	81	86	88	81	82	84
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+	3+	3+	3+	3+	3+	3+	3+	3+	3+
	Without S-9	98	101	100	102	100	100	101	101	99	98	94	97	84	87	89	82	80	85	81	85	87
	Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+	3+	3+	3+	3+	3+	3+	3+	3+	3+



**APPENDIX – 2: INDIVIDUAL PLATE COLONY COUNTS OF REVERTANTS OF TRIAL – I**

Treatment	Test concentration (µg/plate)	Plate No.	No. of Revertants/Plate									
			With S-9					Without S-9				
			TA 98	TA 100	TA 102	TA 1535	TA 1537	TA 98	TA 100	TA 102	TA 1535	TA 1537
Control	100µl of autoclaved distilled water	1	19	101	256	21	8	17	99	250	16	6
		2	18	100	253	20	9	18	101	254	18	5
		3	16	102	236	19	8	16	100	256	20	7
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Vehicle Control	100µl of DMSO	1	17	101	250	20	7	14	102	255	17	8
		2	18	102	256	19	5	17	100	253	18	9
		3	18	99	236	22	9	16	100	260	19	6
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Test item	50	1	16	103	250	20	8	16	101	259	20	5
		2	17	101	252	21	6	15	100	255	18	8
		3	19	100	253	19	7	14	101	256	18	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	100	1	16	101	250	20	6	16	103	260	21	9
		2	16	100	252	21	8	15	100	258	19	7
		3	17	102	251	18	6	14	100	255	16	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	500	1	19	100	256	18	9	15	102	254	16	8
		2	18	98	251	21	5	14	101	258	18	6
		3	17	100	249	16	8	15	100	259	19	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	1000	1	17	101	253	17	7	15	101	255	20	9
		2	16	99	255	16	8	15	102	257	16	6
		3	17	100	240	19	6	15	101	260	17	5
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	2000	1	19	101	251	18	7	15	101	253	19	7
		2	18	99	240	17	5	16	101	257	18	6
		3	18	100	250	17	8	14	100	258	19	7
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Positive Control	100µl	1	196	322	763	198	120	212	315	776	199	171
		2	195	320	765	186	120	214	313	755	194	173
		3	196	321	736	188	121	215	311	780	193	175
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+



**APPENDIX – 3: INDIVIDUAL PLATE COLONY COUNTS OF REVERTANTS OF TRIAL - II**

Treatment	Test concentration (µg/plate)	Plate No.	No. of Revertants/Plate									
			With S-9					Without S-9				
			TA 98	TA 100	TA 102	TA 1535	TA 1537	TA 98	TA 100	TA 102	TA 1535	TA 1537
Control	100µl of autoclaved distilled water	1	17	99	250	16	7	16	99	250	16	10
		2	19	98	254	17	6	19	100	256	17	8
		3	16	99	248	18	8	18	100	254	18	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Vehicle Control	100µl of DMSO	1	19	100	255	19	5	20	101	259	18	10
		2	18	98	257	18	7	19	98	255	16	9
		3	19	98	260	15	6	19	99	254	18	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Test item	50	1	16	101	256	20	9	21	98	257	19	10
		2	19	99	258	17	6	18	98	256	16	10
		3	20	100	259	17	5	19	100	254	19	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	100	1	17	102	256	21	8	16	101	260	20	9
		2	18	101	254	16	5	18	100	257	17	10
		3	16	100	257	16	6	17	98	255	18	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	500	1	18	100	260	15	7	19	102	258	21	7
		2	18	99	258	17	6	17	100	257	18	8
		3	16	99	252	18	6	16	99	252	17	9
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	1000	1	17	101	248	18	6	16	99	260	17	6
		2	16	100	250	19	5	17	99	258	16	8
		3	18	98	251	16	7	17	98	255	18	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
	2000	1	17	98	249	17	8	17	101	253	16	7
		2	16	96	253	15	6	16	100	254	17	9
		3	17	99	255	18	5	17	98	258	17	8
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+
Positive Control	100µl	1	199	320	770	205	120	215	322	769	205	191
		2	192	326	761	198	119	220	327	768	200	202
		3	196	319	763	192	115	218	323	773	195	195
		Lawn Intensity	4+	4+	4+	4+	4+	4+	4+	4+	4+	4+





#### APPENDIX – 4: STERILITY STATUS OF THE STUDY

SI.No.	Sterility Controls	Trial I	Trial II
1	Autoclaved Distilled Water	No microbial contamination	No microbial contamination
2	DMSO	No microbial contamination	No microbial contamination
3	S-9 Mix	No microbial contamination	No microbial contamination
4	Minimal Glucose Agar Plates	No microbial contamination	No microbial contamination
5	Strain Inoculation broth	No microbial contamination	No microbial contamination
6	PBS	No microbial contamination	No microbial contamination
7	Overlay Agar	No microbial contamination	No microbial contamination